Research Paper

Potential Retinal Imaging Markers for Moyamoya Disease: Changes in Retinal Microvasculature and Thickness

Xu Chen, Roel Hubert Louis Haeren, Noël J C Bauer , Rudy M M A Nuijts

University Eye Clinic Maastricht, Maastricht University Medical Center, PO Box 5800, 6202 AZ, Maastricht, The Netherlands

Correspondence to: Rudy M M A Nuijts, email: soraya.jonker@mumc.nlKeywords:Moyamoya disease, Vessel density, retinal thickness, OCTA, Diagnostic markersReceived:May 20, 2023Accepted: July 10, 2024Published: November 9, 2024

Copyright: © 2024 Chen et al. This is an open access article distributed under the terms of the <u>Creative Commons</u> <u>Attribution License (</u>CC BY 3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

ABSTRACT

Purpose: Digital subtraction angiography (DSA) of the brain is the primary means of diagnosing and evaluating Moyamoya disease (MMD). But its high price and invasiveness limit its use as a monitoring tool for disease progression. As a non-invasive test for ophthalmic disorders, the optical coherence tomography angiography (OCTA) is widely used. In addition to ophthalmic diseases, OCTA has also been used in some neurological diseases. The aim of this study was to assess fundus changes in patients with MMD by OCTA and to investigate whether these changes could be a diagnostic and assessment marker for MMD.

Methods: This study evaluated cerebral vessels, superficial macular capillary vessel density (SMC-VD) and macular retinal thickness (MRT) in subjects in the non-operated group (nGO), operated group (OG) and healthy controls (HC) using DSA, OCTA and other techniques. Analyses of variance (ANOVA) and Bonferroni post hoc analysis were used to calculate statistical differences between the three groups. Correlations between SMC-VD and MRT were assessed using Pearson correlation analysis. In addition, the ability of the SMC-VD and the MRT to distinguish MMD from HC was analyzed using receiver operating characteristic (ROC) curves.

Results: We found that the SMC-VD and MRT in the nOG group were significantly lower than those in the HC group and had not returned to normal levels at one month postoperatively. In the nOG, the SMC-VD and MRT were positively correlated in the Full region (6*6 mm) and in the Inner region (3*3 mm), and in many subregions they showed high ability to distinguish MMD from HC. The above findings indicate significant reduction in the SMC-VD and the MRT in patients with MMD even in the absence of ocular clinical manifestation.

Conclusions: SMC-VD and MRT have a strong ability to distinguish between MMD patients and HC, suggesting that OCTA, a relatively inexpensive and non-invasive method, is useful in assessing cerebrovascular changes in MMD patients.

INTRODUCTION

A rare structural abnormality of the cerebral vasculature called Moyamoya disease (MMD). It presents as the proximal intracranial carotid artery (ICA) is stenosed and the distal end is hyperplastic. The abundant collateral vessels at its distal end look like smoke[1; 2; 3; 4]. Recent reports indicate that the average prevalence of MMD is 3.92/100,000 in China and about 10.5/100,000 in Japan, while the prevalence in the United States is only

1/11,000,000. However, in many regions globally, its prevalence has gradually increased in recent years[3; 5; 6; 7]. The main clinical symptom of MMD is cerebral ischemia. Children often present with transient ischemic attack and adults with ischemic cerebral infarction. Complications such as seizures, headache, and cognitive dysfunction may also be associated with cerebral ischemia[2; 3; 8; 9] and cerebral ischemic events have a high risk of recurrence[10]. In previous studies, a few cases of MMD were reported with ocular complications[11; 12; 13; 14]. For example, Ashok et al. reported a young man who had central retinal artery occlusion (CRAO) and was diagnosed with MMD[15]. Alekya et al. reported a case of the patient with both CRAO and MMD, whose optical coherence tomography (OCT) results suggested a reduction in retinal thickness[16].

As a revolutionary imaging method, optical coherence tomography angiography (OCTA) can provide quantitative and morphological data on retinal microvascular changes in vivo[17]. A variety of measures such as blood flow[18], size of FAZ[19], vessel density and others[20] can be calculated using the OCTA technique. In addition, morphological data include vessel length, diameter and number of branches [21]. Currently, OCTA is widely used for the diagnosis and monitoring of ocular diseases which include glaucoma[22], age-related macular degeneration (AMD)[23], and diabetic retinopathy[24]. In the neurological system, OCTA is also considered to assist in monitoring the progression of Alzheimer's disease[25; 26; 27], multiple sclerosis [28; 29; 30], and other diseases.

To investigate whether microvascular and structural changes in the macula reflect the cerebrovascular status of patients with MMD, that is, whether OCTA can be an assistance in the diagnosis of MMD, we collected superficial macular capillary vessel density (SMC-VD) and macular retinal thickness (MRT) in MMD patients and healthy controls (HC). We also collected SMC-VD and MRT data from patients with MMD who had undergone revascularization in a bid to observe the impact of cerebral blood flow alterations on the retina.

MATERIALS AND METHODS

Subject Recruitment

Thirty-two subjects with MMD who met the criteria were recruited by the diagnostic criteria for MMD proposed by the Japanese Committee for the Study of Spontaneous Occlusion of the Ring of Willis[31]. Sixteen of them did not be treated surgically and were the non-operated group (nOG). The other 16 were treated with revascularization (superficial temporal artery-middle cerebral artery anastomosis) and were the operated group (OG). Healthy controls (HC) were included in the study with 16 subjects who met the appropriate criteria. All subjects were assessed between 2018 and 2022 by specialists in the relevant disciplines in the University Eye Clinic Maastricht.

Subjects for inclusion in nOG and OG matched the following criteria: 1) patients with classic MMD diagnosed by Digital subtraction angiography (DSA) and other exams; 2) OG patients undergoing revascularization for one month. Exclusion from the study: 1) DSA results are atypical and/or cannot be confirmed in patients with MMD; 2) other diseases that can affect the vasculature of the brain or a history of brain surgery (e.g. atherosclerosis, congenital malformations of the cerebral vessels, cranio-cerebral tumours, etc.); 3) diseases or history of ocular surgery that can clearly affect the retina and its blood vessels (e.g. glaucoma, ocular tumours. age-related macular degeneration (AMD), etc.); 4) systemic diseases that can affect the retina and its blood vessels (e.g. diabetes, systemic lupus erythematosus (SLE), hypertension, etc.); 5) individuals who are unable to undergo DSA and/or OCT (e.g. with allergic reaction to contrast media).

16 HC were matched for age and gender to nOG and OG patients and met the following criteria: 1) absence of cerebrovascular diseases and history of cranio-cerebral surgery (e.g. MMD, atherosclerosis, cranio-cerebral tumours, etc.); 2) absence of diseases affecting the retina and its blood vessels or history of ocular surgery (e.g. AMD, glaucoma, ocular tumours, etc.); 3) without systemic diseases affecting the retina and its blood vessels (e.g. diabetes, SLE, hypertension, etc.); 4) acceptable for relevant investigations such as DSA and OCTA.

Clinical Examinations

The following medical examinations were performed on all subjects: 1) routine blood and blood biochemical examinations, such as hemoglobin, triglycerides, total cholesterol, glucose, and electrolytes; 2) blood pressure examination; 3) Visual Acuity (VA); 4) DSA of the brain; 5) OCTA. (Some clinical results that did not differ significantly are not shown in the article).

DSA

A DSA machine was applied to the subject to perform DSA of the brain. The patient was advised to take the appropriate position (supine position was chosen for this study). After proper disinfection and anaesthesia, a small incision is opened at the puncture site and a guide wire is inserted using a puncture needle, along which an arterial sheath is placed. The imaging modality (internal carotid arteriography, common carotid arteriography, etc.) is chosen according to the subject's actual condition. A pigtail tube is connected and 5-8 ml of contrast media is injected uniformly at the appropriate location. Frontal and lateral contrast field under fluoroscopy and the results are recorded.

OCTA

We use the Carl Zeiss Meditec AngioPlex system for OCTA imaging. This SD-OCT system operates at an Ascan speed of 68,000 scans per second, a 840 nm light source centre and a 90 nm bandwidth. The system operates with an axial resolution of 5ym, a 15ym lateral resolution and a 2.0mm A-scan depth parameter. Each eye recorded 6*6 mm and 3*3 mm 3D OCTA images. After scanning, each retina of 6*6 mm was divided into nine Early Treatment Diabetic Retinopathy Study (ETDRS) subzones, comprising three concentric circles (1.0 mm, 3.0 mm and 6.0 mm radii, respectively). The 3*3 mm retina was divided into 5 subzones consisting of two concentric circles (radii of 1.0 mm and 3.0 mm, respectively). The vascular density is the percentage of the vascular perfusion area over the measured area. Vessel density was calculated from 2D frontal images created of the superficial retinal layers with a threshold method. Assigned each pixel to perfusion (1) or background (0) by identifying the value of the image block. Results from the region of interest were scaled according to pixel size to calculate the vascular density from the macular centre to the edges of the 6*6 mm and 3*3 mm images. The SMC-VD in this study ranged from the inner limiting membrane (ILM) to the inner plexiform layer (IPL). The left and right eyes of all subjects were evaluated.

Statistical analysis

Analysis of the results was performed with GraphPad Prism version 8 (La Jolla, California, USA) and SPSS version 22.0 (IBM, Armonk, NY, USA). Both ANOVA and Bonferroni Post Hoc Analysis were used for comparison between groups. Linear correlations between MRT and SMC-VD were performed for each group by applying person correlation analysis. For the analysis of differences between HC and MMD patients, subject work characteristic (ROC) curves were plotted.

RESULTS

Fundamental Information

As shown in Table 1, there were 16 subjects in the HC group, nOG and OG, respectively, in this study. It was analyzed for both eyes of each subject. The three groups of subjects did not differ in age (p=0.200), gender (p=0.055) and visual acuity (0.144) between the groups. Figure 1 illustrates typical DSA, SMC-VD and MRT images in MMD patients.

Analysis of the 6*6 mm SMC-VD

SMC-VD in the HC, nOG and the OG in the 6*6 mm region are shown in Table 2 and Figures 2. The SMC-VD was markedly lower in the nOG than HC group in the IS (p=0.020), OS (p=0.001), IN (p=0.016), ON (p<0.001), OI (p=0.028), Outer (p=0.022) and Full (p=0.042) regions. In regions II (p=1.000), IT (p=1.000), OT (p=0.168), C (p=1.000) and Inner (Inner, p=0.134) decreases were not statistically significant (Table 2 and Figure 2a, 2c). In the OS (p=0.036), IN (p=0.040) and ON (p<0.001) subregions, the SMC-VD declined markedly in the OG. The alterations in the remaining subregions were not statistically significant (Table 2 and Figure 2b, 2d). The change in FAZ (p=0.334) was also not statistically significant.

Analysis of the 3*3 mm SMC-VD

SMC-VD changes in the three groups in the 3*3 mm region are shown in Table 3 and Figures 3. In comparison to the HC group, the SMC-VD was markedly decreased in the S (p=0.001), N (p<0.001), I (p=0.042), T (p=0.001) and Inner (p=0.002) regions in nOG patients. Changes in SMC-VD in the C (p=0.084) and Full (p=0.080) regions were not statistically significant (Table 3 and Figure 3a, 3c). SMC-VD in the OG remained markedly lower than in the HC in the S (p=0.026), N (p=0.002), I (p=0.039), C (p=0.018) and Inner (p=0.020) regions. The SMC-VD was not statistically different from HC in T (p=0.686) and Full (p=0.464) subregions of the OG (Table 3 and Figure 3b, 3d). The change in FAZ (p=0.808) was not statistically significant.

MRT

The MRT in the three groups are shown in Table 4 and Figures 4. In all subregions except C (p=0.954), MRT in the nOG was significantly thinner than in the HC group (IT, p=0.011; p<0.001 for all remaining regions; Table 4 and Figure 4a, 4c). Reductions in MRT remained in the IS (p=0.012), IN (p<0.001), ON (p<0.001), Outer (p=0.008) and Full (p=0.023) regions in the OG compared with the HC group (Table 4 and Figure 4b, 4d).

Correlation Analysis

In the nOG, SMC-VD in the Full region was positively correlated with the corresponding MRT value (Pearson r=0.590, p<0.001) (Figure 5a), suggesting that a decrease in MRT was associated with a decrease in SMC-VD. The SMC-VD (3*3 mm) in the Inner region was correlated with the MRT in the corresponding subregion (Figure 5b).

ROC Curves

The ability of MRT to discriminate MMD is represented by the area under the curve (AUC) shown in Figure 6a. The regions with AUC>0.9 were ON (AUC=0.989; p<0.001), OI (AUC=0.909; p<0.001), OT (AUC=0.912; p<0.001), and Outer (AUC=0.969; p<0.001). The analysis of ROC curves for other regions is shown in Table 5. The diagnostic accuracy of SMC-VD for MMD is indicated by the AUC shown in Figure 6b. The only region with AUC>0.9 was ON (AUC=0.907; p<0.001). The analysis of ROC curves for other regions is shown in Table 6.

DISCUSSION

We collected and compared SMC-VD and MRT in the nOG, OG and HC groups to investigate whether microvasculature and retinal thickness were altered in patients with MMD and to assess their diagnostic potential for MMD.

Most subregions of the nOG had lower SMC-VD than HC based on OCTA results (Table 2, 3 and Figures 2, 3). This finding suggests that cerebral vascular disease in patients with MMD may affect superficial retinal capillary density. Several branches are given off by the ICA in the neck and skull, including the ophthalmic artery, which supplies blood to the eyes. The central retinal artery is derived from the ophthalmic artery and supplies blood to the inner five layers of the retina. Blood to the outer five layers of the retina is supplied primarily by the ciliary artery system emanating from the ophthalmic artery [32; 33; 34; 35; 36]. Blood from the central retinal artery supplies the retinal nerve fiber layer, ganglion cell layer and inner plexiform layer through the superficial capillary plexus (SCP), and continues to extend to become the deep capillary plexus (DCP) in the inner nuclear layer. The radial peripapillary capillary plexus (RPCP) is in the nerve fibre layer (NFL)that runs parallel to the axons of the nerve fibres [32; 37; 38; 39; 40]. The capillaries of the SMC-VD we measured in this study are part of the temporal branch of the SCP (Figure 7). The temporal retinal arteries arch around the macula, forming the superior macular arteriole and the arteriola macularis inferior[34; 35] (Figure 7). Stenosis of the ICA is a feature of MMD, and in these cases the blood supply to the branches of the ICA will be reduced. Previous studies have also suggested that retinal ischemia is associated with severe stenosis of the ICA[41; 42]. Therefore, we suggested that the narrowing of ICA led to the reduction of SMC-VD.

Table 1. Fundamental information on the HC, nOG and OG.

	НС	nOG	OG	ANOVA
N	16	16	16	N/A
Eyes (N)	32	32	32	N/A
Age (year)	44.00 ± 12.90	49.44±6.13	49.38±8.71	0.200
Sex, male %	25.00%	18.75%	56.25%	0.055
Disease duration(month)	N/A	14.75 ± 30.40	27.56 ± 19.90	0.169
Visual Acuity (log MAR)	4.67 ± 0.16	4.72 ± 0.17	4.75 ± 0.15	0.144
Systolic blood pressure (mm Hg)	122.86±3.98	121.00±13.87	127.44±18.16	0.384
Diastolic blood pressure (mm Hg)	82.81±4.81	82.82±10.83	84.31±12.24	0.884

ANOVA Showed the Statistical Differences Across Groups. Age, disease duration, visual Acuity and blood puressure were expressed as mean \pm SD.

HC, healthy controls; nOG, non-operated group; OG, operated group; SD, standard deviation.

Subregion	НС	nOG	OG	Bonferroni HC vs. nOG	Bonferroni HC vs. OG	ANOVA
Inner Superior	0.455±0.016	0.438±0.030	0.448±0.029	0.020	0.683	0.025
Outer Superior	0.482±0.012	0.464±0.025	0.470±0.024	0.001#	0.036#	0.002
Inner Nasal	0.449±0.015	0.421±0.053	0.427±0.047	0.016#	0.040#	0.020
Outer Nasal	0.503±0.015	0.469±0.032	0.485±0.014	<0.001#	<0.001#	< 0.001
Inner Inferior	0.444±0.017	0.440±0.023	0.435±0.040	1.000	0.674	0.475
Outer Inferior	0.474±0.018	0.459±0.023	0.467±0.028	0.028	0.765	0.032
Inner Temporal	0.439±0.027	0.433±0.027	0.434±0.038	1.000	1.000	0.654
Outer Temporal	0.445±0.028	0.429±0.039	0.448±0.030	0.168#	0.854#	0.047
Center	0.207 ± 0.060	$0.194{\pm}0.050$	0.178 ± 0.060	1.000	0.130	0.128
Inner	0.446±0.016	0.433±0.025	0.435±0.036	0.130	0.304	0.100
Outer	0.470 ± 0.014	0.456±0.025	$0.467 {\pm} 0.020$	0.020#	0.739#	0.017
Full	0.460±0.013	0.446±0.021	0.454±0.023	0.019	0.663	0.023
FAZ	0.358 ± 0.140	0.311 ± 0.099	$0.358{\pm}0.054$	0.586	1.000	0.334

Table 2. 6*6 mm SMC-VD (mean ± SD) in HC group, nOG and OG

Statistical differences between the nGO, OG and HC groups were calculated by ANOVA and Bonferroni post hoc analysis. #Games-Howell Post Hoc Analysis display the Statistical Differences of the values with the heterogeneous variance.

SMC-VD, superficial macular capillary vessel density; SD, standard deviation; HC, healthy controls; nOG, non-operated group; OG, operated group; FAZ, fluorescein angiography zone.

Subregion	НС	nOG	OG	Bonferroni HC vs. nOG	Bonferroni HC vs. OG	ANOVA	
Superior	0.419 ± 0.022	0.397 ± 0.029	0.403 ± 0.021	0.001	0.026	0.001	
Nasal	0.420 ± 0.016	$0.396 {\pm} 0.032$	$0.397 {\pm} 0.028$	< 0.001	0.002	< 0.001	
Inferior	0.405 ± 0.017	0.388 ± 0.034	0.391 ± 0.025	0.042#	0.039#	0.030	
Temporal	0.410 ± 0.013	$0.388 {\pm} 0.027$	0.405 ± 0.028	0.001#	0.686#	0.001	
Center	$0.187 {\pm} 0.051$	0.158 ± 0.043	0.151 ± 0.060	0.084	0.018	0.015	
Inner	0.412 ± 0.011	0.392 ± 0.028	0.400 ± 0.021	0.002#	0.020#	0.001	
Full	0.379 ± 0.017	0.367 ± 0.027	0.371 ± 0.021	0.080	0.464	0.080	
FAZ	0.360 ± 0.144	0.340 ± 0.099	0.356 ± 0.140	1.000	1.000	0.808	

Table 3. 3*3 mm SMC-VD (mean ± SD) in HC group, nOG and OG

Statistical differences between the nGO, OG and HC groups were calculated by ANOVA and Bonferroni post hoc analysis. # Games-Howell Post Hoc Analysis display the Statistical Differences of the values with the heterogeneous variance.

SMC-VD, superficial macular capillary vessel density; HC, healthy controls; nOG, non-operated group; OG, operated group; FAZ, fluorescein angiography zone.



Fig.1 Typical DSA, SMC-VD and MRT images of MMD patients and HC. (a) and (b) Anteroposterior view and lateral view of normal ICA angiraphy of HC. (c) and (d) Internal carotid angiography in a patient with MMD shows abnormal smoke-like vessels. (e) and (f) OCTA showed SMC-VD of 6*6 mm and 3*3 mm size in HC. (f) and (h) OCTA showing SMC-VD of 6*6 mm and 3*3 mm size in nOG. (i) MRT of HC. (g) MRT of nOG. Compared to HC, patients with MMD showed a significant reduction in SMC-VD and MRT.

DSA, digital subtraction angiography; SMC-VD, superficial macular capillary vessel density; MRT, Macular retinal thickness; MMD, moyamoya disease; nOG, non-operated group; HC, healthy controls; OCTA, Optical Coherence Tomography Angiography.



Fig.2 SMC-VD analysis of 6*6 mm size in HC group, nOG and OG. (a) Statistical analysis of SMC-VD changes in the HC and nOG. IS(p=0.020), OS(p=0.001), IN(p=0.016), ON(p<0.001), OI(p=0.028), Outer(p=0.022) and Full(p=0.042) subregions showed a significant reduction in SMC-VD compared to HC. (b) Statistical analysis of SMC-VD changes in the HC and OG. The SMC-VD in the OS(p=0.036), IN(p=0.040) and ON(p<0.001) subregions were significantly lower than that in the HC group. (c) Mean values of SMC-VD (6*6 mm) in each subregion of HC group and nOG. The red area indicates a significant decrease in SMC-VD in this region compared to HC (d) Mean values of SMC-VD (6*6 mm) in each subregion of HC group and OG. The red area indicates a significant decrease in SMC-VD in this region compared to HC (d) Mean values of SMC-VD (6*6 mm) in each subregion of HC group and OG. The red area indicates a significant decrease in SMC-VD in this region compared to HC (d) Mean values of SMC-VD (6*6 mm) in each subregion of HC group and OG. The red area indicates a significant decrease in SMC-VD in this region compared to HC (d) Mean values of SMC-VD (6*6 mm) in each subregion of HC group and OG. The red area indicates a significant decrease in SMC-VD in this region compared to HC. * p<0.05; ** p<0.01; *** p<0.001.

SMC-VD, superficial macular capillary vessel density; HC, healthy controls; nOG, non-operated group; OG, operated group; IS, inner superior; OS, outer superior; IN, inner nasal; ON, outer nasal; OI, outer inferior; FAZ, fluorescein angiography zone.



Fig.3 SMC-VD analysis of 3*3 mm size in HC group, nOG and OG. (a) Statistical analysis of SMC-VD changes in the HC and nOG. S(p=0.001), N(p<0.001), I(p=0.042), T(p=0.001) and Inner(p=0.002) subregions showed a significant reduction in SMC-VD compared to HC. (b) Statistical analysis of SMC-VD changes in the HC and OG. SMC-VD was significantly lower in the S (p=0.026), N (p=0.002), I (p=0.039), C (p=0.018) and Inner (p=0.020) subregions than in the HC group. (c) Mean values of SMC-VD (3*3 mm) in each subregion of HC group and nOG. The red area indicates a significant decrease in SMC-VD in this region compared to HC. (d) Mean values of SMC-VD (3*3 mm) in each subregion of HC group and OG. The red area indicates a significant decrease in SMC-VD in this region compared to HC. (d) mean values of SMC-VD (3*3 mm) in each subregion of HC group and OG. The red area indicates a significant decrease in SMC-VD in this region compared to HC. (d) Mean values of SMC-VD (3*3 mm) in each subregion of HC group and OG. The red area indicates a significant decrease in SMC-VD in this region compared to HC. (d) mean values of SMC-VD (3*3 mm) in each subregion of HC group and OG. The red area indicates a significant decrease in SMC-VD in this region compared to HC. * p<0.05; ** p<0.01; *** p<0.001.

SMC-VD, superficial macular capillary vessel density; HC, healthy controls; nOG, non-operated group; OG, operated group; S, superior; N, nasal; I, inferior; T, temporal; C, center; FAZ, fluorescein angiography zone.

These changes suggest questions about their role in structural and functional variations in the retina. In the present study, MRT of patients in the nOG was significantly thinner than that of healthy subjects in all subregions but C (Table 4 and Figures 4), yet these subjects did not show significant visual impairment. Blockage of the central retinal artery and its branches can often be observed as retinal edema [33; 34; 39; 43; 44]. Due to ischemia and hypoxia, the retinal cells become edematous, and if ischemia is prolonged the inner retinal cells gradually become necrotic, leading to reduced retinal thickness [34; 39]. The MMD subjects we included in our study all had long disease duration (Table 1). Therefore, we interpret their MRT reduction as an alteration caused by prolonged ischemia and hypoxia. That means the reduction

of MRT is associated with the reduction of SMC-VD. And a correlation was found in nOG between MRT and SMC-VT in the Full region at 6*6mm (Figure 5a) and in the Inner region at 3*3mm (Figure 5b). This evidence indicates a close association between reduced MRT and reduced SMC-VD in patients with MMD.

To further investigate this link in patients with MMD we evaluated structural and microvascular variations in the retinas of MMD subjects one month after revascularization. The results indicated that in this group (OG) the SMC-VD remained significantly lower than HC in OS, IN, ON, S, I and Inner regions (Table 2, 3 and Figure 2b, 2d, 3b, 3d). The MRT did not return to normal in the IS, IN, ON, Outer, and Full regions compared to HC (Table 4 and Figure 4b, 4d). We speculate that the above results may be due to the occurrence of ischemia-reperfusion injury. Ischemia-reperfusion injury mainly refers to tissue damage caused by the restoration of blood supply after ischemia or/and

hypoxia[45; 46]. This can manifest as endothelial damage, increased permeability and leukocyte obstruction in microvessels, especially small

Subregion	НС	nOG	OG	Bonferroni	Bonferroni	ANOVA	
				HC vs. nOG	HC vs. OG		
Inner Superior	332.44±14.70	312.94±15.15	320.69±17.94	<0.001	0.012	< 0.001	
Outer Superior	298.59±20.72	277.81±16.06	287.91±18.23	<0.001	0.068	< 0.001	
Inner Nasal	330.06±15.11	311.00±15.35	307.19±19.40	<0.001	< 0.001	< 0.001	
Outer Nasal	317.25±8.14	290.69±11.24	274.28±13.85	< 0.001	< 0.001	< 0.001	
Inner Inferior	323.50±14.32	304.41±22.22	321.88±19.58	< 0.001	1.000	< 0.001	
Outer Inferior	279.19±5.38	262.50±13.37	282.00±14.87	<0.001#	0.578#	< 0.001	
Inner temporal	316.50±14.28	303.25±14.96	315.84±22.88	0.011	1.000	0.005	
Outer Temporal	275.69±7.12	261.13±9.17	295.34±22.38	<0.001#	<0.001#	< 0.001	
Center	238.75±22.85	243.91±21.78	241.78±16.40	0.954	1.000	0.603	
Inner	325.75±12.83	308.03±14.30	316.47±18.74	< 0.001	0.056	< 0.001	
Outer	292.75±7.44	273.19±11.27	$285.00{\pm}10.94$	< 0.001	0.008	< 0.001	
Full	301.28±9.49	285.31±10.06	294.19±11.50	< 0.001	0.023	< 0.001	

Table 4. MRT (mean ± SD) in HC group, nOG and OG

Statistical differences between the nGO, OG and HC groups were calculated by ANOVA and Bonferroni post hoc analysis. # Games-Howell Post Hoc Analysis display the Statistical Differences of the values with the heterogeneous variance.MRT, macular retinal thickness; HC, healthy controls; nOG, non-operated group; OG, operated group; FAZ, fluorescein angiography zone.



Fig.4 MRT analysis of HC group, nOG and OG. (a) Statistical analysis of MRT changes in the HC and nOG. All subregions except subregion C displayed a marked reduction in MRT compared to HC (IT, p=0.011; p<0.001 for all remaining regions). (b) Statistical analysis of MRT changes in the HC and OG. The MRT was markedly thinner in the IS (p=0.012), IN (p<0.001), ON (p<0.001), OT (p<0.001), Outer (p=0.008) and Full (p=0.023) regions in comparison to the HC group. (c) Mean values of MRT in each subregion of HC group and nOG. The red area indicates a significant decrease in MRT in this region compared to HC. (d) Mean values of MRT in each subregion of HC group and OG. The red area indicates a significant decrease in MRT in this region compared to HC * p<0.05; ** p<0.01; *** p<0.001.

MRT, macular retinal thickness; HC, healthy controls; nOG, non-operated group; OG, operated group; IS, inner superior; OS, outer superior; IN, inner nasal; ON, outer nasal; II, inner inferior; OI, outer inferior; IT, inner temporal; OT, outer temporal; C, center; FAZ, fluorescein angiography zone.



Fig.5 The correlation between MRT and SMC-VD. (a) Full SMC-VD of 6*6mm sizes in MMD patients (nOG) was positively correlated with the corresponding MRT (Person r=0.590, p<0.001). (b) In patients with MMD (nOG), a correlation was found between the full SMC-VD of 3 x 3 mm and the corresponding MRT values (Person r=0.434, p=0.013). MRT, Macular retinal thickness; SMC-VD, superficial macular capillary vessel density; MMD, moyamoya disease; HC, healthy controls; nOG, non-operated group.

arteries and capillaries[47; 48]. A recent case report also showed that a patient with MMD who underwent left superficial temporal artery-middle cerebral artery anastomosis for one month had a significantly reduced superficial retinal vessels of the left eye which did not return to normal until nine months postoperatively. This vascular complex, however, did not show any significant abnormality before surgery [49]. It has also been shown that patients with MMD have cerebral hyperperfusion and a range of complications after revascularization treatment [50; 51]. On this basis, it seems feasible that in the present study small arteries and capillaries in the macula may have suffered reperfusion injury in addition to ischemia. In future studies we may need more specialized neurologists to evaluate the results of cerebral revascularization, rather than assuming that the cerebral vessels are back to normal just by the results of DSA.

Although the cerebral vascular system has a complex structure, Goto et al. suggested that retinal vasculopathy may in part reflect cerebral microvascular lesions[52; 53; 54]. In embryology, the retina was considered to be the central nervous system's extension with a common origin in the embryonic neural tube. The anterior part of the neural tube differentiates into brain tissue and forms a pair of optic vesicles which develop into the right and left retinas[35; 55; 56]. Moreover, anatomically, blood of the retina, the anterior two-thirds of the telencephalon, and part of the mesencephalon are supplied by the ICA[35; 36; 57] and alterations in the retinal arteries are related to vascular disease in the brain [58; 59]. Strokes

have been linked to retinal vessel changes even since the early studies using fundus photography[60; 61] and changes in cognitive abilities[62; 63]. We therefore hypothesized that OCTA measurement of retinal vascular structure and perfusion may also reflect in part the cerebral blood flow in patients with MMD. We evaluated the diagnostic accuracy of MRT and SMC-VD for MMD using ROC curves. The results showed high AUC values for MRT in all sub-regions except for region C. The AUC values of ON, OI, OT and Outer were all over 0.9, and approaching this value (at 0.87) in the Full region (Table 5 and Figure 6a). It is suggested that MRT has strong potential to discriminate between MMD and HC. However, for SMC-VD only ON had an AUC greater than 0.9, values of the remaining subregions being mostly in the range of 0.6-0.7 (Table 6 and Figure 6b). The diagnostic accuracy of SMC-VD for MMD was therefore inferior to that of MRT. In our analysis, this may reflect the fact that vascular lesions in MMD affect not only the superficial retinal vessels, but also the deeper vascular layers, such as those of the choroid.

However, the limitations of our study need to be considered. For example, our sample size was so small that it limited the accuracy of the study. In the study, we only focused on the capillaries in the superficial layers of the retina. Other vessels of the retina, such as the DCP and choroidal capillaries, may also altered. And, we should assess the correspondence between cerebral and retinal vascular alterations by long-term, relatively frequent DSA and OCTA in future studies.

Table 5. ROC Curve Analysis of WIRT						
MRT	AUC	SE	95% CI	P value		
Inner	0.815	0.052	0.714 ± 0.017			
Superior			0.714 to 0.917	< 0.001		
Outer	0.839	0.050	0.741 to 0.937			
Superior				< 0.001		
Inner Nasal	0.796	0.054	0.690 to 0.902	< 0.001		
Outer Nasal	0.989	0.009	0.972 to 1.000	< 0.001		
Inner Inferior	0.804	0.056	0.695 to 0.913	< 0.001		
Outer Inferior	0.909	0.039	0.832 to 0.986	< 0.001		
Inner	0.735	0.062	0.614 to 0.856	0.001		
Temporal						
Outer	0.912	0.034	0.845 to 0.979			
Temporal				< 0.001		
Center	0.574	0.072	0.432 to 0.716	0.308		

Inner	0.818	0.051	0.718 to 0.919	< 0.001
Outer	0.969	0.018	0.933 to 1.000	< 0.001
Full	0.872	0.042	0.790 to 0.955	< 0.001

AUC values, SE, 95% CI and P-values for each subregion.

ROC, receiver operating characteristic; MRT, Macular retinal thickness; AUC, area under ROC curve; SE, standard errors; 95% CI, 95% confidence intervals.

SMC-VD(6*6mm)	AUC	SE	95% CI	P value
Inner Superior	0.672	0.067	0.540 to 0.804	0.018
Outer Superior	0.731	0.062	0.609 to 0.852	0.002
Inner Nasal	0.648	0.069	0.513 to 0.784	0.041
Outer Nasal	0.907	0.038	0.832 to 0.981	< 0.001
Inner Inferior	0.560	0.075	0.413 to 0.706	0.413
Outer Inferior	0.698	0.067	0.567 to 0.829	0.007
Inner Temporal	0.598	0.073	0.454 to 0.742	0.177
Outer Temporal	0.607	0.072	0.466 to 0.748	0.142
Center	0.600	0.072	0.459 to 0.741	0.171
Inner	0.648	0.069	0.513 to 0.783	0.042
Outer	0.651	0.069	0.516 to 0.786	0.038
Full	0.673	0.067	0.543 to 0.804	0.017

Table 6. ROC Curve Analysis of SMC-VD(6*6mm)

AUC values, SE, 95% CI and P-values for each subregion.

ROC, receiver operating characteristic; SMC-VD, superficial macular capillary vessel density;

AUC, area under ROC curve; SE, standard errors; 95% CI, 95% confidence intervals.



Fig.6 ROC curves of MRT and SMC-VD. (a) The AUC suggests the accuracy of MRT for the diagnosis of MMD. The AUC in the ON (AUC=0.989), OI(AUC=0.909), OT(AUC=0.912) and Outer (AUC=0.969) subregions was greater than 0.9. (b) The AUC suggests accuracy of SMC-VD for MMD diagnosis. The AUC in the ON (AUC= 0.907) subregion was greater than 0.9.

ROC, receiver operating characteristic; MRT, Macular retinal thickness; SMC-VD, superficial macular capillary vessel density; AUC, area under ROC curve; MMD, moyamoya disease; ON, outer nasal; OI, outer inferior; OT, outer temporal.



Fig.7 Anatomical localization of retinal vessels and anatomical stratification of the retina. (a) The area circled by the red square is the approximate location of the macula as described in the study. (b)The SMC-VD we focused on in the study was mainly from superficial capillary plexus.

SMC-VD, superficial macular capillary vessel density.

CONCLUSION

In summary, we conclude that patients with MMD still have significant SMC-VD and MRT reduction even in the absence of ocular clinical manifestations. And MRT reduction is closely associated with SMC-VD reduction. These reductions are not immediately restored with the reestablishment of cerebral blood flow. This may be related to ischemia-reperfusion damage to the retinal arteries. Importantly, the SMC-VD and MRT have a strong ability to distinguish MMD from HC, suggesting that alterations in retinal microvasculature and structure may be a cheaper non-invasive marker for assessing vascular changes in the brain in patients with MMD. The OCTA may serve as a more convenient way to monitor changes in MMD.

AUTHOR CONTRIBUTIONS

(I) Conception and design: RN; (II) Administrative support: NB; (III) Provision of study materials or patients: All authors; (IV) Collection and assembly of data: XC; (V) Data analysis and interpretation: RH; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

CONFLICT OF INTERESTS

There is no declaration for authors about conflict of

interests in this study.

Data Availability Statements: The data that support the findings of this study are available from the corresponding author upon reasonable request.

Open Access Statement: In accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), this article is distributed as Open Access. Under this license, the article may be reproduced and distributed for non-commercial purposes only if no changes or edits have been made and the original work is properly cited (Links to official publications are available by relevant DOI and license).

FUNDING

No funding.

REFERENCES

- J. Suzuki, and A. Takaku, Cerebrovascular "moyamoya" disease. Disease showing abnormal net-like vessels in base of brain. Arch Neurol 20 (1969) 288-99.
- R.M. Scott, and E.R. Smith, Moyamoya disease and moyamoya syndrome. N Engl J Med 360 (2009) 1226-37.
- S. Shang, D. Zhou, J. Ya, S. Li, Q. Yang, Y. Ding, X. Ji, and R. Meng, Progress in moyamoya disease. Neurosurg Rev 43 (2020) 371-382.

- T.S. Kiyoshi Sato, Moyamoya Disease, Cerebrovascular Diseases in Children, Springer, New York, NY, 1992, pp. 227-243.
- Y. Goto, and Y. Yonekawa, Worldwide distribution of moyamoya disease. Neurol Med Chir (Tokyo) 32 (1992) 883-6.
- S. Kuroda, and K. Houkin, Moyamoya disease: current concepts and future perspectives. Lancet Neurol 7 (2008) 1056-66.
- T. Hishikawa, K. Sugiu, and I. Date, Moyamoya Disease: A Review of Clinical Research. Acta Med Okayama 70 (2016) 229-36.
- S.K. Kim, B.K. Cho, J.H. Phi, J.Y. Lee, J.H. Chae, K.J. Kim, Y.S. Hwang, I.O. Kim, D.S. Lee, J. Lee, and K.C. Wang, Pediatric moyamoya disease: An analysis of 410 consecutive cases. Ann Neurol 68 (2010) 92-101.
- K. Kazumata, K.K. Tha, H. Narita, I. Kusumi, H. Shichinohe, M. Ito, N. Nakayama, and K. Houkin, Chronic ischemia alters brain microstructural integrity and cognitive performance in adult moyamoya disease. Stroke 46 (2015) 354-60.
- D. Chiu, P. Shedden, P. Bratina, and J.C. Grotta, Clinical features of moyamoya disease in the United States. Stroke 29 (1998) 1347-51.
- T.L. Slamovits, T.G. Klingele, R.M. Burde, and M.H. Gado, Moyamoya disease with central retinal vein occlusion. Case report. J Clin Neuroophthalmol 1 (1981) 123-7.
- R. Chace, and T.R. Hedges, 3rd, Retinal artery occlusion due to moyamoya disease. J Clin Neuroophthalmol 4 (1984) 31-4.
- S. Ushimura, K. Mochizuki, M. Ohashi, S. Ito, and H. Hosokawa, Sudden blindness in the fourth month of pregnancy led to diagnosis of moyamoya disease. Ophthalmologica 207 (1993) 169-73.
- P. Albrecht, C. Blasberg, S. Lukas, M. Ringelstein, A.K. Müller, J. Harmel, E.M. Kadas, D. Finis, R. Guthoff, O. Aktas, H.P. Hartung, F. Paul, A.U. Brandt, P. Berlit, A. Methner, and M. Kraemer, Retinal pathology in idiopathic moyamoya angiopathy detected by optical coherence tomography. Neurology 85 (2015) 521-7.
- 15. M.A. Kumar, and B.A. Ganesh, CRAO in Moyamoya Disease. J Clin Diagn Res 7 (2013) 545-7.
- 16. A.P. Rajanala, H.T. Le, and M.K. Gill, Central retinal

artery occlusion as initial presentation of Moyamoya disease in a middle-aged woman. Am J Ophthalmol Case Rep 18 (2020) 100705.

- J.P. Campbell, M. Zhang, T.S. Hwang, S.T. Bailey, D.J. Wilson, Y. Jia, and D. Huang, Detailed Vascular Anatomy of the Human Retina by Projection-Resolved Optical Coherence Tomography Angiography. Sci Rep 7 (2017) 42201.
- R.K. Wang, Q. Zhang, Y. Li, and S. Song, Optical coherence tomography angiography-based capillary velocimetry. J Biomed Opt 22 (2017) 66008.
- J. Choi, J. Kwon, J.W. Shin, J. Lee, S. Lee, and M.S. Kook, Quantitative optical coherence tomography angiography of macular vascular structure and foveal avascular zone in glaucoma. PLoS One 12 (2017) e0184948.
- H. Akil, K.G. Falavarjani, S.R. Sadda, and A.A. Sadun, Optical Coherence Tomography Angiography of the Optic Disc; an Overview. J Ophthalmic Vis Res 12 (2017) 98-105.
- A. Wylęgała, F. Wylęgała, and E. Wylęgała, Aflibercept Treatment Leads to Vascular Abnormalization of the Choroidal Neovascularization. J Healthc Eng 2018 (2018) 8595278.
- N.K. Scripsema, P.M. Garcia, R.D. Bavier, T.Y. Chui, B.D. Krawitz, S. Mo, S.A. Agemy, L. Xu, Y.B. Lin, J.F. Panarelli, P.A. Sidoti, J.C. Tsai, and R.B. Rosen, Optical Coherence Tomography Angiography Analysis of Perfused Peripapillary Capillaries in Primary Open-Angle Glaucoma and Normal-Tension Glaucoma. Invest Ophthalmol Vis Sci 57 (2016) Oct611-oct620.
- L. Roisman, and R. Goldhardt, OCT Angiography: An Upcoming Non-invasive Tool for Diagnosis of Agerelated Macular Degeneration. Curr Ophthalmol Rep 5 (2017) 136-140.
- K.B. Schaal, M.R. Munk, I. Wyssmueller, L.E. Berger, M.S. Zinkernagel, and S. Wolf, VASCULAR ABNORMALITIES IN DIABETIC RETINOPATHY ASSESSED WITH SWEPT-SOURCE OPTICAL COHERENCE TOMOGRAPHY ANGIOGRAPHY WIDEFIELD IMAGING. Retina 39 (2019) 79-87.
- 25. L. Ferrari, S.C. Huang, G. Magnani, A. Ambrosi, G. Comi, and L. Leocani, Optical Coherence

Tomography Reveals Retinal Neuroaxonal Thinning in Frontotemporal Dementia as in Alzheimer's Disease. J Alzheimers Dis 56 (2017) 1101-1107.

- C.Y. Santos, L.N. Johnson, Y.Y. Lim, B.M. Fernandez, B.R. Ott, S. Salloway, and P.J. Snyder, P4–058.: Retinal Nerve Fiber Layer and Ganglion Cell Layer Volume Changes in Preclinical Alzheimer's Disease over 27 Months. Alzheimer's & Dementia 13 (2017).
- C.Y. Santos, L.N. Johnson, S.E. Sinoff, E.K. Festa, W.C. Heindel, and P.J. Snyder, Change in retinal structural anatomy during the preclinical stage of Alzheimer's disease. Alzheimers Dement (Amst) 10 (2018) 196-209.
- R. Behbehani, A. Abu Al-Hassan, A. Al-Salahat, D. Sriraman, J.D. Oakley, and R. Alroughani, Optical coherence tomography segmentation analysis in relapsing remitting versus progressive multiple sclerosis. PLoS One 12 (2017) e0172120.
- N. Feucht, M. Maier, G. Lepennetier, M. Pettenkofer, C. Wetzlmair, T. Daltrozzo, P. Scherm, C. Zimmer, M.M. Hoshi, B. Hemmer, T. Korn, and B. Knier, Optical coherence tomography angiography indicates associations of the retinal vascular network and disease activity in multiple sclerosis. Mult Scler 25 (2019) 224-234.
- F.C. Oertel, H.G. Zimmermann, A.U. Brandt, and F. Paul, Novel uses of retinal imaging with optical coherence tomography in multiple sclerosis. Expert Rev Neurother 19 (2019) 31-43.
- M. Fujimura, and T. Tominaga, Diagnosis of moyamoya disease: international standard and regional differences. Neurol Med Chir (Tokyo) 55 (2015) 189-93.
- J. Stone, D. van Driel, K. Valter, S. Rees, and J. Provis, The locations of mitochondria in mammalian photoreceptors: relation to retinal vasculature. Brain Res 1189 (2008) 58-69.
- B. Mac Grory, M. Schrag, V. Biousse, K.L. Furie, M. Gerhard-Herman, P.J. Lavin, L. Sobrin, S.I. Tjoumakaris, C.M. Weyand, and S. Yaghi, Management of Central Retinal Artery Occlusion: A Scientific Statement From the American Heart Association. Stroke 52 (2021) e282-e294.
- 34. K.B.F.D.S.W.F.M.L.A. Yannuzzi, The Retinal Atlas, second deition, 2017.
- 35. Z.D. Shitian Zhong, Zengtao Wang, Zuguo Liu,

Jianhua Yan, Clinical Anatomy of Ophthalmology, second edition, 2020.

- N.W. Zuguo Liu, Xinghuai Sun, Luosheng Tang 眼科 学基础, 人民卫生出版社, 2018.
- P. Henkind, Radial peripapillary capillaries of the retina. I. Anatomy: human and comparative. Br J Ophthalmol 51 (1967) 115-23.
- 38. J.M. Provis, Development of the primate retinal vasculature. Prog Retin Eye Res 20 (2001) 799-821.
- 39. T.W.G. Antonia M. Joussen, Bernd Kirchhof, Stephen J, Ryan, Retinal Vascular Disease, 2011.
- 40. P.R. David R.Chow, OCT Angiography, 2018.
- 41. J.B. Mizener, P. Podhajsky, and S.S. Hayreh, Ocular ischemic syndrome. Ophthalmology 104 (1997) 859-64.
- V. Babikian, C.A. Wijman, B. Koleini, S.N. Malik, N. Goyal, and I.C. Matjucha, Retinal ischemia and embolism. Etiologies and outcomes based on a prospective study. Cerebrovasc Dis 12 (2001) 108-13.
- 43. S.S. Hayreh, Acute retinal arterial occlusive disorders. Prog Retin Eye Res 30 (2011) 359-94.
- G.A. Ochakovski, D.A. Wenzel, M.S. Spitzer, S. Poli, F. Härtig, M.D. Fischer, S. Dimopoulos, and M. Schultheiss, Retinal oedema in central retinal artery occlusion develops as a function of time. Acta Ophthalmol 98 (2020) e680-e684.
- R.Q. Jianzhi Wang, Liling Wu, Liankun Sun, Wenbin Li, Zhisheng Jiang 病理生理学, 人民卫生出版社, 2018.
- 46. Y.G. Xianzhong Xiao, 病理生理学, 高等教育出版社, 2018.
- D.L. Carden, and D.N. Granger, Pathophysiology of ischaemia–reperfusion injury. The Journal of Pathology 190 (2000) 255-266.
- 48. H.K. Eltzschig, and T. Eckle, Ischemia and reperfusion--from mechanism to translation. Nat Med 17 (2011) 1391-401.
- K.J. Lo, A.G. Wang, and H.C. Cheng, Retinal Vasculature Changes after Bypass Surgery for Moyamoya Disease. Ophthalmology 128 (2021) 120.
- 50. S. Kuroda, H. Kamiyama, H. Abe, K. Asaoka, and K. Mitsumori, Temporary neurological deterioration caused by hyperperfusion after extracranial-intracranial bypass--case report and study of

cerebral hemodynamics. Neurol Med Chir (Tokyo) 34 (1994) 15-9.

- 51. M. Fujimura, T. Kaneta, S. Mugikura, H. Shimizu, and T. Tominaga, Temporary neurologic deterioration due to cerebral hyperperfusion after superficial temporal artery-middle cerebral artery anastomosis in patients with adult-onset moyamoya disease. Surg Neurol 67 (2007) 273-82.
- I. Goto, S. Katsuki, H. Ikui, K. Kimoto, and T. Mimatsu, Pathological studies on the intracerebral and retinal arteries in cerebrovascular and noncerebrovascular diseases. Stroke 6 (1975) 263-9.
- 53. N. Patton, T. Aslam, T. Macgillivray, A. Pattie, I.J. Deary, and B. Dhillon, Retinal vascular image analysis as a potential screening tool for cerebrovascular disease: a rationale based on homology between cerebral and retinal microvasculatures. J Anat 206 (2005) 319-48.
- H.E. Moss, Retinal Vascular Changes are a Marker for Cerebral Vascular Diseases. Curr Neurol Neurosci Rep 15 (2015) 40.
- D.L. Stenkamp, Development of the Vertebrate Eye and Retina. Prog Mol Biol Transl Sci 134 (2015) 397-414.
- J.B. Miesfeld, and N.L. Brown, Eye organogenesis: A hierarchical view of ocular development. Curr Top Dev Biol 132 (2019) 351-393.
- T.H. Rim, A.W.J. Teo, H.H.S. Yang, C.Y. Cheung, and T.Y. Wong, Retinal Vascular Signs and Cerebrovascular Diseases. J Neuroophthalmol 40 (2020) 44-59.
- 58. A.D. Hughes, E. Falaschetti, N. Witt, S. Wijetunge,

S.A. Thom, T. Tillin, S.J. Aldington, and N. Chaturvedi, Association of Retinopathy and Retinal Microvascular Abnormalities With Stroke and Cerebrovascular Disease. Stroke 47 (2016) 2862-2864.

- 59. P. Nelis, I. Kleffner, M.C. Burg, C.R. Clemens, M. Alnawaiseh, J. Motte, M. Marziniak, N. Eter, and F. Alten, OCT-Angiography reveals reduced vessel density in the deep retinal plexus of CADASIL patients. Sci Rep 8 (2018) 8148.
- R.I. Lindley, J.J. Wang, M.C. Wong, P. Mitchell, G. Liew, P. Hand, J. Wardlaw, D.A. De Silva, M. Baker, E. Rochtchina, C. Chen, G.J. Hankey, H.M. Chang, V.S. Fung, L. Gomes, and T.Y. Wong, Retinal microvasculature in acute lacunar stroke: a crosssectional study. Lancet Neurol 8 (2009) 628-34.
- R. Kawasaki, M.Z. Che Azemin, D.K. Kumar, A.G. Tan,
 G. Liew, T.Y. Wong, P. Mitchell, and J.J. Wang,
 Fractal dimension of the retinal vasculature and
 risk of stroke: a nested case-control study.
 Neurology 76 (2011) 1766-7.
- N. Patton, A. Pattie, T. MacGillivray, T. Aslam, B. Dhillon, A. Gow, J.M. Starr, L.J. Whalley, and I.J. Deary, The association between retinal vascular network geometry and cognitive ability in an elderly population. Invest Ophthalmol Vis Sci 48 (2007) 1995-2000.
- J. Ding, M.W. Strachan, F.G. Fowkes, T.Y. Wong, T.J. Macgillivray, N. Patton, T.A. Gardiner, I.J. Deary, and J.F. Price, Association of retinal arteriolar dilatation with lower verbal memory: the Edinburgh Type 2 Diabetes Study. Diabetologia 54 (2011) 1653-62.